

Product Name: Recombinant Schistosoma Japonicum GST
Catalog #: PEV0739




Summary

Name	Glutathione S-transferase Class-mu 26 kDa Isozyme/GST
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Schistosoma Japonicum Glutathione S-transferase Class-mu 26 kDa Isozyme is produced by our E.coli expression system and the target gene encoding Met1-Lys218 is expressed.
Accession #	P08515
Host	E.coli
Species	Schistosoma Japonicum
Predicted Molecular Mass	25.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Schistosoma Japonicum GST
Catalog #: PEV0739



Alternative Names

Glutathione S-transferase class-mu 26 kDa isozyme; GST 26; Sj26 antigen; SjGST

Background

Glutathione S-transferases (GSTs), previously known as ligandins, comprise a family of eukaryotic and prokaryotic phase II metabolic isozymes best known for their ability to catalyze the conjugation of the reduced form of glutathione (GSH) to xenobiotic substrates for the purpose of detoxification. The GST family consists of three superfamilies: the cytosolic, mitochondrial, and microsomal (MAPEG) proteins. GST isoenzymes appear to play a central role in the parasite detoxification system. Other functions are also suspected including a role in increasing the solubility of haematin in the parasite gut. The activity of GSTs is dependent upon a steady supply of GSH from the synthetic enzymes gamma-glutamylcysteine synthetase and glutathione synthetase, as well as the action of specific transporters to remove conjugates of GSH from the cell. The primary role of GSTs is to detoxify xenobiotics by catalyzing the nucleophilic attack by GSH on electrophilic carbon, sulfur, or nitrogen atoms of said nonpolar xenobiotic substrates, thereby preventing their interaction with crucial cellular proteins and nucleic acids.

Note

For Research Use Only , Not for Diagnostic Use.