Product Name: Recombinant Rabbit TNF alpha

Catalog #: PEV1680



Summary

Name TNF alpha/TNFSF2/TNFα

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Rabbit Tumor Necrosis Factor Alpha is produced by our E.coli

expression system and the target gene encoding Val78-Leu235 is expressed.

Accession # P04924

Host E.coli

Species Rabbit

Predicted Molecular Mass 17.59 KDa

Formulation Lyophilized from a 0.2 μm filtered solution of 20mM PB, 250mM NaCl, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Lyophilized protein should be stored at \leq -20°C, stable for one year after receipt.

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at \leq -20°C for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than $100\mu g/ml$. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than $100\mu g/ml$. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Rabbit TNF alpha

Catalog #: PEV1680





Alternative Names

Tumor Necrosis Factor; Cachectin; TNF-Alpha; Tumor Necrosis Factor Ligand Superfamily Member 2; TNF-a; TNF; TNFA; TNFSF2

Background

Tumor necrosis factor alpha (TNF α) is the prototypic ligand of the TNF superfamily. TNF α forms a homotrimer and functions by activating two types of receptors TNF-R1 (TNF receptor type 1,p55R) and TNF-R2 (TNF receptor type 2,p75R). TNFα is a pleiotropic cytokine that is capable to promote inflammation, to induce apoptotic cell death, and to inhibit tumorigenesis and viral replication. TNF α is a potent lymphoid factor that exerts cytotoxic effects on a wide range of tumor cells and certain other target cells.

Note

For Research Use Only, Not for Diagnostic Use.