


Product Name: Recombinant Rhesus Macaque IL-33 (N-6His)
Catalog #: PEV2058



Summary

| | |
|---------------------------------|--|
| Name | IL-33/Interleukin-33/IL-1F11/Interleukin-1 Family Member 11/Nuclear Factor From High Endothelial Venules/NF-HEV/C9orf26/NFHEV |
| Purity | Greater than 95% as determined by reducing SDS-PAGE |
| Endotoxin level | <1 EU/μg as determined by LAL test. |
| Construction | Recombinant Rhesus Macaque Interleukin 33 is produced by our E.coli expression system and the target gene encoding Ser112-Ile270 is expressed with a 6His tag at the N-terminus. |
| Accession # | F7DLG4 |
| Host | E.coli |
| Species | Rhesus macaque |
| Predicted Molecular Mass | 20.3 KDa |
| Formulation | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| Stability&Storage | Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles. |
| Reconstitution | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

SDS-PAGE image



Alternative Names



Product Name: Recombinant Rhesus Macaque IL-33 (N-6His)
Catalog #: PEV2058

Interleukin-33; IL-33; Interleukin-1 Family Member 11; IL-1F11; Nuclear Factor From High Endothelial Venules; NF-HEV; IL33; C9orf26; IL1F11; NFHEV

Background

Interleukin-33 (IL-33) belongs to the IL-1 superfamily. IL33 is highly expressed in endothelial venules found in tonsils, Peyer patches and mesenteric lymph nodes, but almost undetectable in placenta. IL33 induces the production of T helper-2 (Th2)-associated cytokines. IL-33 is a cytokine that mediates its biological effects by binding to and signals through IL1RL1/ST2 and IL-1 Receptor Accessory Protein (IL1RAP), activating intracellular molecules in the NF- κ B and MAP kinase signaling pathways that drive production of type 2 cytokines from polarized Th2 cells.

Note

For Research Use Only , Not for Diagnostic Use.