

Product Name: Recombinant Human Acrp30 (C-6His)
Catalog #: PHH0022




Summary

Name	Adiponectin/Acrp30/ADIPOQ
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Adipocyte Complement-Related 30 kDa Protein is produced by our Mammalian expression system and the target gene encoding Glu19-Asn244 is expressed with a 6His tag at the C-terminus.
Accession #	Q15848
Host	Human Cells
Species	Human
Predicted Molecular Mass	25.58 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human Acrp30 (C-6His)
Catalog #: PHH0022



Alternative Names

Adiponectin; 30 kDa Adipocyte Complement-Related Protein; Adipocyte complement-related 30 kDa protein; ACRP30; Adipocyte; C1q and Collagen Domain-Containing Protein; Adipose Most Abundant Gene Transcript 1 Protein; apM-1; Gelatin-Binding Protein; ADIPOQ; ACDC; ACRP30; APM1; GBP28

Background

Adiponectin is a secreted protein. It is synthesized exclusively by adipocytes and secreted into plasma. Adiponectin is an important adipokine that is involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Adiponectin Stimulates AMPK phosphorylation and activates in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Adiponectin also antagonizes TNF-alpha by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. It inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. Adiponectin may play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex: LMW, MMW or HMW.

Note

For Research Use Only , Not for Diagnostic Use.