

## Summary

Name	Alcohol dehydrogenase class 4 mu/sigma chain/ADH7
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/ $\mu$ g as determined by LAL test.
Construction	Recombinant Human Alcohol Dehydrogenase Class 4 Mu/Sigma Chain is produced by our Mammalian expression system and the target gene encoding Met1-Phe386 is expressed with a 6His tag at the C-terminus.
Accession #	P40394
Host	Human Cells
Species	Human
Predicted Molecular Mass	42.5 KDa
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at $\leq$ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $\leq$ -20°C for 3 months
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image





## Background

Alternative NamesAlcohol Dehydrogenase Class 4 Mu/Sigma Chain; Alcohol Dehydrogenase Class IV<br/>Mu/Sigma Chain; Gastric Alcohol Dehydrogenase; Retinol Dehydrogenase; ADH7BackgroundAlcohol dehydrogenase class 4 mu/sigma chain (ADH7) is a cytoplasm enzyme<br/>which is a member of the alcohol dehydrogenase family. The expression of this<br/>gene makes it much more abundant in the stomach than the liver, thus it differs<br/>from the other known gene family members. ADH7 may participate in the<br/>synthesis of retinoic acid, a hormone important for cellular differentiation.<br/>Medium-chain (octanol) and aromatic (m-nitrobenzaldehyde) compounds are the<br/>best substrates. Ethanol is not a good substrate but at the high ethanol<br/>concentrations reached in the digestive tract, it plays a role in the ethanol<br/>oxidation and contributes to the first pass ethanol metabolism.

## Note

For Research Use Only, Not for Diagnostic Use.