## **Product Name: Recombinant Human AGP2 (C-6His)**

Catalog #: PHH0046



### **Summary**

Name alpha-1-acid glycoprotein 2/AGP2/ORM2

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/µg as determined by LAL test.

Construction Recombinant Human Alpha-1-acid Glycoprotein 2 is produced by our

Mammalian expression system and the target gene encoding Gln19-Ser201 is

expressed with a 6His tag at the C-terminus.

Accession # P19652

Host **Human Cells** 

**Species** Human

**Predicted Molecular Mass** 22.6 KDa

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. **Formulation** 

**Shipping** The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Lyophilized protein should be stored at  $\leq$  -20°C, stable for one year after receipt. Stability&Storage


Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at  $\leq$  -20°C for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### **SDS-PAGE** image



## **Background**

# **Product Name: Recombinant Human AGP2 (C-6His)**

Catalog #: PHH0046



Alternative Names Orosomucoid-2; OMD 2; ORM2; AGP2

Background Alpha-1-acid glycoprotein 2 is a secreted protein that belongs to the calycin

superfamily, lipocalin family. It is expressed by the liver and secreted in plasma. It appears to function in modulating the activity of the immune system during the acute-phase reaction. It functions as transport protein in the blood stream. It binds various hydrophobic ligands in the interior of its beta-barrel domain. It also binds

synthetic drugs and influences their distribution and availability.

#### Note

For Research Use Only , Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838