Product Name: Recombinant Human ApoM (C-6His)

Catalog #: PHH0092



Summary

Name Apolipoprotein M/ApoM

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Apolipoprotein M is produced by our Mammalian

expression system and the target gene encoding Met1-Asn188 is expressed

with a 6His tag at the C-terminus.

Accession # 095445

Host Human Cells

Species Human

Predicted Molecular Mass 22.29 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Lyophilized protein should be stored at \leq -20°C, stable for one year after receipt.


Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at \leq -20°C for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human ApoM (C-6His) Catalog #: PHH0092



Alternative Names Apolipoprotein M; Apo-M; ApoM; Protein G3a; APOM; G3A; NG20

Background Apolipoprotein M is a secreted protein which belongs to the Lipocalin family.

ApoM often presents in high density lipoprotein (HDL) and to a lesser extent in triglyceride-rich lipoproteins (TGRLP) and low density lipoproteins (LDL). The ApoM gene encoded protein is expressed in liver and kidney, secreted through the plasma membrane but remains membrane-bound. ApoM probably involved in lipid transport. ApoM can bind sphingosine-1-phosphate, myristic acid, palmitic acid and stearic acid, retinol, all-trans-retinoic acid and 9-cis-retinoic acid. The expression of ApoM could be regulated by platelet activating factor (PAF), Transforming Growth Factors (TGF), Insulin-Like Growth factor (IGF) and Leptin.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838