Product Name: Recombinant Human BTN3A3 (C-6His) Catalog #: PHH0179



Summary

Name BTN3A3/Butyrophilin subfamily 3 member A3

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Butyrophilin Subfamily 3 Member A3 is produced by

our Mammalian expression system and the target gene encoding Gln30-

Trp248 is expressed with a 6His tag at the C-terminus.

Accession # 000478

Host Human Cells

Species Human

Predicted Molecular Mass 24.6 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human BTN3A3 (C-6His)

Catalog #: PHH0179





Alternative Names

Butyrophilin subfamily 3 member A3;BTN3A3

Background

Human BTN3A3, also known as butyrophilin subfamily 3 member A3 and BTF3, is a Single-pass type I membrane protein which belongs to the immunoglobulin superfamily and BTN/MOG family. The butyrophilin (BTN) genes are a group of major histocompatibility complex (MHC)-associated genes that encode type I membrane proteins with 2 extracellular immunoglobulin domains and an intracellular B30.2 (PRYSPRY) domain. It can be detected in peripheral blood mononuclear cells, T-cells I, spleen and lymphocytes. BTN3A3 plays a role in T-cell responses in the adaptive immune response.

Note

For Research Use Only, Not for Diagnostic Use.