

Product Name: Recombinant Human CDH17 (C-6His)
Catalog #: PHH0192




Summary

Name	Cadherin-17/CDH17
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Cadherin-17 is produced by our Mammalian expression system and the target gene encoding Gln23-Met787 is expressed with a 6His tag at the C-terminus.
Accession #	AAI13465.1
Host	Human Cells
Species	Human
Predicted Molecular Mass	86 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human CDH17 (C-6His)
Catalog #: PHH0192



Alternative Names

Cadherin-17; Intestinal Peptide-Associated Transporter HPT-1; Liver-Intestine Cadherin; LI-Cadherin; CDH17

Background

Cadherin-17 is a single-pass type I membrane protein that belongs to the cadherin superfamily. Cadherin-17 consists of one extracellular region containing seven cadherin domains and one transmembrane region but it lacks the conserved cytoplasmic domain. Cadherin-17 is expressed in the gastrointestinal tract and pancreatic duct. Cadherins are calcium dependent cell adhesion proteins and preferentially interact with each other in a homophilic manner in connecting cells. Cadherin-17 may have a role in the morphological organization of liver and intestine and involved in intestinal peptide transport.

Note

For Research Use Only , Not for Diagnostic Use.