


**Product Name: Recombinant Human TF (C-6His)**  
**Catalog #: PHH0292**



## Summary

<b>Name</b>	Coagulation Factor III/Tissue Factor/CD142/F3/TF
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Coagulation Factor III/Tissue Factor is produced by our Mammalian expression system and the target gene encoding Gly34-Glu251 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	P13726
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	25.76 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human TF (C-6His)**  
**Catalog #: PHH0292**

---



**Alternative Names**

Tissue Factor; TF; Coagulation Factor III; Thromboplastin; CD142; F3

**Background**

Tissue Factor (TF) is a single-pass type I membrane glycoprotein member of the tissue factor family. TF expression is highly dependent upon cell type. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. TF initiates blood coagulation by forming a complex with circulating factor VII or VIIa. The complex activates factors IX or X by specific limited proteolysis. TF plays a role in normal hemostasis by initiating the cell-surface assembly and propagation of the coagulation protease cascade.

**Note**

For Research Use Only , Not for Diagnostic Use.