Product Name: Recombinant Human IL-17RA (C-Fc)

Catalog #: PHH0311



Summary

Name IL-17RA/CD217/IL-17 R alpha/IL-17 receptor A

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Interleukin-17 Receptor A is produced by our

Mammalian expression system and the target gene encoding Leu33-Trp320 is

expressed with a human IgG1 Fc tag at the C-terminus.

Accession # Q96F46

Host Human Cells

Species Human

Predicted Molecular Mass 60.6 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is


not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human IL-17RA (C-Fc)

Catalog #: PHH0311





Alternative Names

Interleukin-17 receptor A; IL-17 receptor A; IL-17RA; CDw217; CD_antigen; CD217; IL17RA; IL17R

Background

Interleukin 17A (IL17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A. Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors, this receptor likely has a multimeric structure.

Note

For Research Use Only, Not for Diagnostic Use.