# **Summary**

Name TRAIL R3/CD263/TNFRSF10C/TRID

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Human TNF-Related Apoptosis-Inducing Ligand Receptor 3 is

produced by our Mammalian expression system and the target gene encoding Ala26-Ala221 is expressed with a human IgG1 Fc, 6His tag at the C-

terminus.

Accession # O14798

**Host** Human Cells

**Species** Human

Predicted Molecular Mass 48.7 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw


cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838



### **Alternative Names**

Tumor Necrosis Factor Receptor Superfamily Member 10C; Antagonist Decoy Receptor for TRAIL/Apo-2L; Decoy TRAIL Receptor Without Death Domain; Decoy Receptor 1; DcR1; Lymphocyte Inhibitor of TRAIL; TNF-Related Apoptosis-Inducing Ligand Receptor 3; TRAIL Receptor 3; TRAIL-R3; TRAIL Receptor Without an Intracellular Domain; CD263; TNFRSF10C; DCR1; LIT; TRAILR3; TRID

# **Background**

Tumor Necrosis Factor Receptor Superfamily Member 10C (TNFRSF10C) is a glycosyl-phosphatidylinositol-linked membrane protein which binds TRAIL with high affinity. TNFRSF10C has the TRAIL-binding extracellular cysteine-rich domains, lacks the intracellular signaling domain. As a result, binding of TRAIL to TRAIL R3 doesn't transduce an apoptosis signal. The expression of TRAIL R3 gene has been shown to protect cells bearing TRAIL R1 and/or TRAIL R2 from TRAIL-induced apoptosis.

#### Note

For Research Use Only, Not for Diagnostic Use.