

Product Name: Recombinant Human CLEC10A (C-6His)
Catalog #: PHH0326




Summary

Name	CLEC10A/CD301
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human C-Type Lectin Domain Family 10 Member A is produced by our Mammalian expression system and the target gene encoding Gln61-His316 is expressed with a 6His tag at the C-terminus.
Accession #	Q8IUN9
Host	Human Cells
Species	Human
Predicted Molecular Mass	29.8 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human CLEC10A (C-6His)
Catalog #: PHH0326



Alternative Names

C-Type Lectin Domain Family 10 Member A; C-Type Lectin Superfamily Member 14; Macrophage Lectin 2; CD301; CLEC10A; CLECSF13; CLECSF14; HML

Background

C-Type Lectin Domain Family 10 Member A (CLEC10A) is a type II transmembrane C-type lectin that is expressed on immature myeloid dendritic cells and alternatively activated (tolerogenic) macrophages. CLEC10A/MGL binds and internalizes molecules with terminal nonsialylated GalNAc carbohydrates such as the Tn carcinoma antigen. CLEC10A/MGL also binds the GP envelope glycoprotein on Marburg and Ebola viruses and enhances viral entry and infectivity. It constitute a unique class of C-type lectins because of their specificity for galactose and its structural homologues. CLEC10A is thought to participate in the recognition of molecules from both altered self and pathogens due to its monosaccharide specificity for Gal and N-acetylgalactosamine (GalNAc). Human and rat carry a single gene for CLEC10A/MGL, while mouse has two closely related MGL1 and MGL2 genes.

Note

For Research Use Only , Not for Diagnostic Use.