


**Product Name: Recombinant Human CD32b (C-6His)**  
**Catalog #: PHH0330**



## Summary

<b>Name</b>	Fc gamma RIIB/CD32b/Low affinity immunoglobulin gamma Fc region receptor II
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Low Affinity Immunoglobulin Gamma Fc Region Receptor II-B is produced by our Mammalian expression system and the target gene encoding Thr43-Pro217 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	P31994
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	20.64 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human CD32b (C-6His)**  
**Catalog #: PHH0330**

---



**Alternative Names**

Low Affinity Immunoglobulin Gamma Fc Region Receptor II-b; IgG Fc Receptor II-b; CDw32; Fc-Gamma RII-b; Fc-Gamma-RIIb; FcRII-b; CD32; FCGR2B; FCG2; IGFR2

**Background**

FcγRIIB is a low affinity receptor that recognizes the Fc portion of IgG. The human CD32 group consists of FcγRIIA, FcγRIIB, and FcγRIIC proteins that share 94-99% sequence identity in their extracellular domains but differ substantially in their transmembrane and cytoplasmic domains. FcγRII protein is expressed on cells of both myeloid and lymphoid lineages as well as on cells of non-hematopoietic origin. FcγRIIB has an intrinsic cytoplasmic immunoreceptor tyrosine-based inhibitory motif (ITIM) and delivers an inhibitory signal upon ligand binding. Ligation of FcγRIIB on B cells down-regulates antibody production and in some circumstances may promote apoptosis. Co-ligation of FcγRIIB on dendritic cells inhibits maturation and blocks cell activation. FcγRIIB may also be a target for monoclonal antibody therapy for malignancies. FcγRIIB plays an important negative-regulating role through modulating the signals from activation receptors.

**Note**

For Research Use Only , Not for Diagnostic Use.