


Product Name: Recombinant Human ICAM-1 (C-Fc)
Catalog #: PHH0358



Summary

Name	ICAM-1/ICAM1/CD54/Intercellular adhesion molecule 1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Intercellular Adhesion Molecule 1 is produced by our Mammalian expression system and the target gene encoding Asn26-Glu480 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	P05362
Host	Human Cells
Species	Human
Predicted Molecular Mass	76.8 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human ICAM-1 (C-Fc)
Catalog #: PHH0358



Alternative Names

Intercellular Adhesion Molecule 1; ICAM-1; Major Group Rhinovirus Receptor; CD54; ICAM1

Background

Inter-Cellular Adhesion Molecule 1 (ICAM1) is a type of intercellular adhesion molecule continuously present in low concentrations in the membranes of leukocytes and endothelial cells. As an endothelial and leukocyte-associated transmembrane protein, ICAM1 is well known for its importance in stabilizing cell-cell interactions and facilitating leukocyte endothelial transmigration. The presence of heavy glycosylation and other structural characteristics lend ICAM1 binding sites for a number of immune-associated ligands. Notably, ICAM-1 binds to macrophage adhesion ligand-1 (Mac-1; ITGB2 / ITGAM), leukocyte function associated antigen-1 (LFA-1/integrin), and fibrinogen. ICAM-1 expressed by respiratory epithelial cells is also the binding site for rhinovirus, the causative agent of most common colds.

Note

For Research Use Only , Not for Diagnostic Use.