Summary

Name CD66b/CEACAM8

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Human Carcinoembryonic Antigen-Related Cell Adhesion

> Molecule 8 is produced by our Mammalian expression system and the target gene encoding Gln35-His141 is expressed with a 6His tag at the C-terminus.

Accession # P31997

Host **Human Cells**

Species Human

Predicted Molecular Mass 13.03 KDa

Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. **Formulation**

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 Stability&Storage


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human CEACAM8 (C-6His) Enkilife Catalog #: PHH0363

Alternative Names Carcinoembryonic Antigen-Related Cell Adhesion Molecule 8; CD67 Antigen;

Carcinoembryonic Antigen CGM6; Non-Specific Cross-Reacting Antigen NCA-95;

CD66b; CEACAM8; CGM6

BackgroundCarcinoembryonic Antigen-Related Cell Adhesion Molecule 8 (CEACAM8) is a single chain, GPI-anchored, highly glycosylated protein which belongs to the

single chain, GPI-anchored, highly glycosylated protein which belongs to the immunoglobulin superfamily and the carcinoembryonic antigen(CEA) family. CEACAM8 is expressed by neutrophils and eosinophils, and serves as a binding partner for CEACAM-6 and Galectin-3. It contains two Ig-like C2-type (immunoglobulin-like) domains and one Ig-like V-type (immunoglobulin-like) domain. Mature human CEACAM8 is a 287 amino acid GPI-linked glycoprotein. CEACAM family members are a set of widely expressed proteins involved in several biological functions, including cell adhesion, migration, signal transduction, and the regulation of gene expression. Abnormal overexpression and downregulation

of some CEACAMs have been described in tumor cells.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838