

Product Name: Recombinant Human CEACAM3 (C-6His)
Catalog #: PHH0364




Summary

Name	CD66d/CEACAM3
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Carcinoembryonic Antigen-Related Cell Adhesion Molecule 3 is produced by our Mammalian expression system and the target gene encoding Lys35-Gly155 is expressed with a 6His tag at the C-terminus.
Accession #	P40198
Host	Human Cells
Species	Human
Predicted Molecular Mass	14.13 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human CEACAM3 (C-6His)
Catalog #: PHH0364



Alternative Names

Carcinoembryonic Antigen-Related Cell Adhesion Molecule 3; Carcinoembryonic Antigen CGM1; CD66d; CEACAM3; CD66D; CGM1

Background

Carcinoembryonic Antigen-Related Cell Adhesion Molecule 3 (CEACAM3) belongs to the immunoglobulin superfamily and CEA family. CEACAM3 was originally described in bile ducts of liver as biliary glycoprotein. Subsequently, it was found to be a cell-cell adhesion molecule detected on leukocytes, epithelia, and endothelia. CEACAM3 mediates cell adhesion via homophilic as well as heterophilic binding to other proteins of the subgroup. In addition, it is associated with the differentiation and arrangement of tissue three-dimensional structure, angiogenesis, apoptosis, tumor suppression, metastasis, and the modulation of innate and adaptive immune responses.

Note

For Research Use Only , Not for Diagnostic Use.