


Product Name: Recombinant Human FAS (C-6His)
Catalog #: PHH0380



Summary

Name	CD95/TNFRSF6/FAS
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human TNF Receptor Superfamily Member 6 is produced by our Mammalian expression system and the target gene encoding Gln26-Asn173 is expressed with a 6His tag at the C-terminus.
Accession #	P25445
Host	Human Cells
Species	Human
Predicted Molecular Mass	17.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human FAS (C-6His)
Catalog #: PHH0380



Background

Alternative Names

Tumor necrosis factor receptor superfamily member 6; Apo-1 antigen; Apoptosis-mediating surface antigen FAS; FASLG receptor; APT1; FAS1; TNFRSF6 and FAS

Background

FAS is a receptor and contains three TNFR-Cys repeats and one death domain. It has been shown that FAS is involved in the physiological regulation of programmed cell death, and has been implicated in the pathogenesis of various malignancies and diseases of the immune system. FADD (adapter molecule) recruits caspase-8 to the activated receptor, the resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases mediating apoptosis. FAS-mediated apoptosis may play a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both.

Note

For Research Use Only , Not for Diagnostic Use.