


## Summary

<b>Name</b>	CEACAM5/CEACAM-5/CD66e/CEA
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Carcinoembryonic Antigen-Related Cell Adhesion Molecule 5 is produced by our Mammalian expression system and the target gene encoding Lys35-Ala685 is expressed with a human IgG1 Fc tag at the C-terminus.
<b>Accession #</b>	NP_004354.3
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	98.5 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human CEACAM5 (C-Fc)**  
**Catalog #: PHH0394**



---

**Alternative Names**

Carcinoembryonic antigen-related cell adhesion molecule 5; CEACAM5; Carcinoembryonic antigen; CEA; Meconium antigen 100; CD66e

**Background**

Carcinoembryonic antigen-related cell adhesion molecules (CEACAMs) belong to a group of mammalian immunoglobulin related glycoproteins. They play critical roles in cell–cell recognition. CEACAM5, also called CEA and CD66e, is characterized by having seven extracellular Ig domains and a glycosylphosphatidylinositol (GPI) anchor. CEACAM5 is expressed primarily by epithelial cells, and functions as a calcium-independent adhesion molecule through homophilic and heterophilic interactions with CEACAM1. Studies have shown that CEACAM5 is overexpressed in a majority of carcinomas, and its overexpression can protect tumor cells from apoptosis. It is commonly used as a cancer marker.

**Note**

For Research Use Only , Not for Diagnostic Use.