

**Product Name: Recombinant Human Cryptic (C-6His)**  
**Catalog #: PHH0455**




## Summary

|                                 |  |
|---------------------------------|--|
| <b>Name</b>                     | Cryptic protein/cryptic  |
| <b>Purity</b>                   | Greater than 95% as determined by reducing SDS-PAGE  |
| <b>Endotoxin level</b>          | <1 EU/μg as determined by LAL test.  |
| <b>Construction</b>             | Recombinant Human Cryptic Protein is produced by our Mammalian expression system and the target gene encoding Tyr26-Gly169 is expressed with a 6His tag at the C-terminus.   |
| <b>Accession #</b>              | P0CG37   |
| <b>Host</b>                     | Human Cells  |
| <b>Species</b>                  | Human  |
| <b>Predicted Molecular Mass</b> | 16.9 KDa   |
| <b>Formulation</b>              | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.  |
| <b>Shipping</b>                 | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.   |
| <b>Stability&amp;Storage</b>    | Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.  |
| <b>Reconstitution</b>           | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

## SDS-PAGE image

**Product Name: Recombinant Human Cryptic (C-6His)**  
**Catalog #: PHH0455**



### **Alternative Names**

Cryptic protein;Cryptic family protein 1;CFC1

### **Background**

Cryptic (CFC1) is a member of the epidermal growth factor (EGF)- Cripto, Frl-1, and Cryptic (CFC) family. It contains an EGF-like domain, and is glycosylated on its N-terminal during post-translational modification. Cryptic is identified as a NODAL coreceptor involved in the correct establishment of the left-right axis. It may play a role in mesoderm and/or neural patterning during gastrulation. The unnormal expression of this gene may causes a series of diseases such as HTX2, Transposition of the great arteries dextro-looped 2, and Conotruncal heart malformations.

### **Note**

For Research Use Only , Not for Diagnostic Use.