Summary

Name CXCL10/IP-10/CRG-2/INP10/SCYB10

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human C-X-C Motif Chemokine 10 is produced by our

Mammalian expression system and the target gene encoding Val22-Pro98 is

expressed with a human IgG1 Fc, 6His tag at the C-terminus.

Accession # P02778

Host Human Cells

Species Human

Predicted Molecular Mass 36.6 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Lyophilized protein should be stored at \leq -20°C, stable for one year after receipt.


Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at \leq -20°C for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human CXCL10 (C-Fc-6His) Enkilife Catalog #: PHH0466

Alternative Names C-X-C Motif Chemokine 10; 10 kDa Interferon Gamma-Induced Protein; Gamma-

IP10; IP-10; Small-Inducible Cytokine B10; CXCL10; INP10; SCYB10

Background Human C-X-C Motif Chemokine Ligand 10 (CXCL10) is a non-ELR chemokine

secreted by various cell types, such as monocytes, endothelial cells and fibroblasts, in response to IFN-γ. CXCL10 functions via chemokine receptor CXCR3. CXCL10 has been attributed to several roles, such as chemoattraction for activated T-

lymphocytes, inhibition of angiogenesis, and antitumor activity.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838