


Product Name: Recombinant Human CDK2AP2 (C-6His)
Catalog #: PHH0484



Summary

Name	CDK2AP2/Cyclin-dependent kinase 2-associated protein 2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Cyclin-Dependent Kinase 2-Associated Protein 2 is produced by our Mammalian expression system and the target gene encoding Met1-Thr126 is expressed with a 6His tag at the C-terminus.
Accession #	O75956
Host	Human Cells
Species	Human
Predicted Molecular Mass	14.1 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human CDK2AP2 (C-6His)
Catalog #: PHH0484



Alternative Names

Cyclin-dependent kinase 2-associated protein 2;CDK2-associated protein 2;DOC-1-related protein;DOC-1R;CDK2AP2;DOC1R

Background

CDK2AP2, also known as DOC1R, is short for cyclin-dependent kinase 2-associated protein 2. The gene CDK2AP2 encodes this protein that interacts with cyclin-dependent kinase 2 associated protein 1. Pseudogenes associated with this gene are located on chromosomes 7 and 9. Alternatively spliced transcript variants have been observed for this gene. It belongs to the CDK2AP family. CDK2AP1 (cyclin-dependent kinase 2-associated protein 1), corresponding to the gene doc-1 (deleted in oral cancer 1), is a tumor suppressor protein. The doc-1 gene is absent or down-regulated in hamster oral cancer cells and in many other cancer cell types. The ubiquitously expressed CDK2AP1 protein is the only known specific inhibitor of CDK2, making it an important component of cell cycle regulation during G(1)-to-S phase transition.

Note

For Research Use Only , Not for Diagnostic Use.