

**Product Name: Recombinant Human Cyr61 (C-Fc)**  
**Catalog #: PHH0488**




## Summary

<b>Name</b>	Cyr61/CCN1/Cysteine-rich angiogenic inducer 61
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Cysteine-rich Angiogenic Inducer 61 is produced by our Mammalian expression system and the target gene encoding Thr25-Asp381 is expressed with a human IgG1 Fc tag at the C-terminus.
<b>Accession #</b>	O00622
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	66.5 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 6% Trehalose, 4% Mannitol, 0.05% Tween80, pH7.5.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Human Cyr61 (C-Fc)**  
**Catalog #: PHH0488**



### Alternative Names

Protein CYR61;CCN family member 1; Cysteine-rich angiogenic inducer 61;Insulin-like growth factor-binding protein 10 ; GIG1; CYR61; CCN1;IGFBP10;

### Background

Protein CYR61, also known as CCN family member 1, Cysteine-rich angiogenic inducer 61,Insulin-like growth factor-binding protein 10 , GIG1, CYR61, CCN1 and IGFBP10, belongs to the CCN family, CYR61 is a secreted protein and contains one CTCK (C-terminal cystine knot-like) domain,one IGFBP N-terminal domain,one TSP type-1 domain and one VWFC domain. CYR61 promotes cell proliferation, chemotaxis, angiogenesis and cell adhesion. CYR61 plays important roles in inflammation and tissue repair. CYR61 is associated with diseases related to chronic inflammation, including rheumatoid arthritis, atherosclerosis, diabetes-related nephropathy and retinopathy, and many different forms of cancers.

### Note

For Research Use Only , Not for Diagnostic Use.