## **Product Name: Recombinant Human CST2 (C-6His)**

Catalog #: PHH0501



### **Summary**

Name Cystatin SA/CST2/Cystatin-2

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Human Cystatin SA is produced by our Mammalian expression

system and the target gene encoding Trp21-Ala141 is expressed with a 6His

tag at the C-terminus.

Accession # P09228

**Host** Human Cells

**Species** Human

Predicted Molecular Mass 15.39 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH

8.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image



## **Background**

# Product Name: Recombinant Human CST2 (C-6His) Catalog #: PHH0501



Alternative Names Cystatin-SA; Cystatin-2; Cystatin-S5; CST2

**Background** Cystatin SA is a member of family 2 of the Cystatin superfamily. Together with

Cystatins S and SN, it is produced by the salivary gland and secreted largely in the submandibular/sublingual saliva. Cystatin SA inhibits Cathepsin L, but may not inhibit Cathepsins B and H. The functions of Cystatin SA are largely unknown. However, it has been suggested to play a role in the control of periodontal tissue

destruction.

#### Note

For Research Use Only , Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838