Product Name: Recombinant Human CRISP-3 (C-6His) Catalog #: PHH0503



Summary

Name Cysteine-rich secretory protein 3/CRISP-3

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Cysteine-Rich Secretory Protein 3 is produced by our

Mammalian expression system and the target gene encoding Asn21-Tyr245 is

expressed with a 6His tag at the C-terminus.

Accession # P54108

Host Human Cells

Species Human

Predicted Molecular Mass 26.54 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than $100\mu g/ml$. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than $100\mu g/ml$. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human CRISP-3 (C-6His) Catalog #: PHH0503



Background

Alternative Names Cysteine-Rich Secretory Protein 3; CRISP-3; Specific Granule Protein of 28 kDa;

SGP28; CRISP3

Background Cysteine-rich secretory protein 3 (CRISP-3) is a secreted protein, containing 1 SCP

domain and 1 ShKT domain.It is belongs to the CRISP family. CRISP-3 is a glycoprotein that belongs to the family of cysteine-rich secretory proteins (CRISPs) which was originally discovered in human neutrophilic granulocytes. CRISP-3 is also widely distibuted in exocrine glands (salivary glands, pancreas and prostate), eosinophilic granulocytes and to a lower level in epididymis, ovary, thymus and colon. The presence of CRISP-3 in neutrophils, eosinophils and in exocrine secretions indicates a role in innate host defense. The antibody has been raised against recombinant C-terminally truncated form of CRISP-3 and recognizes both

the N-glycosylated and non-glycosylated form of the mature protein.

Note

For Research Use Only , Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838