

**Product Name: Recombinant Human DKK-1 (N-8His)**  
**Catalog #: PHH0533**




---

## Summary

<b>Name</b>	DKK-1/Dickkopf-related protein 1
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<0.01 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Dickkopf-related Protein 1 is produced by our Mammalian expression system and the target gene encoding Thr32-His266 is expressed with a 8His tag at the N-terminus.
<b>Accession #</b>	O94907
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	27 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Human DKK-1 (N-8His)**  
**Catalog #: PHH0533**



### Alternative Names

Dickkopf-related protein 1; Dickkopf-1; Dkk-1

### Background

Dickkopf-related protein 1 (DKK-1), is a member of the dickkopf family. DKK1 secreted proteins with two cysteine-rich domains separated by a linker region. It antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer disease.

### Note

For Research Use Only , Not for Diagnostic Use.