

Product Name: Recombinant Human EFNA1 (C-Fc)
Catalog #: PHH0587




Summary

Name	Ephrin-A1/EFNA1/TNFAIP4
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Ephrin-A1 is produced by our Mammalian expression system and the target gene encoding Asp19-Ser182 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	AAH32698.1
Host	Human Cells
Species	Human
Predicted Molecular Mass	46.5 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human EFNA1 (C-Fc)
Catalog #: PHH0587



Alternative Names

Ephrin-A1; EPH-Related Receptor Tyrosine Kinase Ligand 1; LERK-1; Immediate Early Response Protein B61; Tumor Necrosis Factor Alpha-Induced Protein 4; TNF Alpha-Induced Protein 4; EFNA1; EPLG1; LERK1; TNFAIP4

Background

Ephrin-A1 is a member of the A-type ephrin family of cell surface proteins that function as ligands for the A-type Eph receptor tyrosine kinase family. Ephrin-A1 can be induced by TNF and IL1B. Its expression levels can be down-regulated in primary glioma tissues compared to the normal tissues. The soluble monomeric form is expressed in the glioblastoma multiforme (GBM) and breast cancer cells. Soluble Ephrin-A1 is necessary for the transformation of HeLa and SK-BR3 cells and participates in the relocalization of EPHA2 away from sites of cell-cell contact during transformation. Ephrin-A1 plays an important role in angiogenesis and tumor neovascularization.

Note

For Research Use Only , Not for Diagnostic Use.