


**Product Name: Recombinant Human EFNA3 (C-6His)**  
**Catalog #: PHH0588**



## Summary

|                                 |  |
|---------------------------------|--|
| <b>Name</b>                     | Ephrin-A3/EFNA3  |
| <b>Purity</b>                   | Greater than 95% as determined by reducing SDS-PAGE  |
| <b>Endotoxin level</b>          | <1 EU/μg as determined by LAL test.  |
| <b>Construction</b>             | Recombinant Human Ephrin-A3 is produced by our Mammalian expression system and the target gene encoding Gln23-Ser211 is expressed with a 6His tag at the C-terminus.   |
| <b>Accession #</b>              | P52797   |
| <b>Host</b>                     | Human Cells  |
| <b>Species</b>                  | Human  |
| <b>Predicted Molecular Mass</b> | 22.25 KDa  |
| <b>Formulation</b>              | Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.  |
| <b>Shipping</b>                 | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.   |
| <b>Stability&amp;Storage</b>    | Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.   |
| <b>Reconstitution</b>           | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

## SDS-PAGE image



## Background

**Product Name: Recombinant Human EFNA3 (C-6His)**  
**Catalog #: PHH0588**

---



**Alternative Names**

Ephrin-A3; EFL-2; EHK1 Ligand; EHK1-L; EPH-Related Receptor Tyrosine Kinase Ligand 3; LERK-3; EFNA3; EFL2; EPLG3; LERK3

**Background**

Ephrins-A3 belongs to the Ephrins ligand family, which is involved in a variety of biological processes, especially in the nervous system and in erythropoiesis. It is shown that Ephrin-A3 is expressed in brain, skeletal muscle, spleen, thymus, prostate, testis, ovary, small intestine, and peripheral blood leukocytes. Ephrin-A3 has a GPI anchor following the extracellular sequence and a signal sequence of 22 amino acids. Ephrin-A3 can bind EphA2, EphA3, EphA4, EphA5, EphA6, EphA7, EphA8 and EphB1. Furthermore, it is associated with tumor growth and metastasis.

**Note**

For Research Use Only , Not for Diagnostic Use.