


Product Name: Recombinant Human Fetuin A (C-6His)
Catalog #: PHH0638



Summary

Name	Fetuin A/AHSG
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Alpha-2-HS-Glycoprotein is produced by our Mammalian expression system and the target gene encoding Ala19-Val367 is expressed with a 6His tag at the C-terminus.
Accession #	AAH48198.1
Host	Human Cells
Species	Human
Predicted Molecular Mass	38.36 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 7.5.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human Fetuin A (C-6His)
Catalog #: PHH0638



Alternative Names

Alpha-2-HS-Glycoprotein; Alpha-2-Z-Globulin; Ba-Alpha-2-Glycoprotein; Fetuin-A; AHSB; FETUA

Background

α -2-HS Glycoprotein (AHSB) is a glycoprotein that is composed of two subunits, the A and B chains, belongs to the Cystatin family of proteases inhibitors. It is highly expressed in embryonic cells and adult hepatocytes, and is expressed to a lesser extent in monocytes/macrophages. AHSB is an important circulating inhibitor of calcification in vivo, and is downregulated during the acute-phase response. It is involved in several functions, such as endocytosis, brain development and the formation of bone tissue. In addition, AHSB may influence the resolution of inflammation by modulating the phagocytosis of apoptotic cells by macrophages. AHSB blocks TGF-beta-dependent signaling in osteoblastic cells.

Note

For Research Use Only , Not for Diagnostic Use.