


Summary

Name	FLRT3
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Fibronectin Leucine Rich Transmembrane Protein 3 is produced by our Mammalian expression system and the target gene encoding Lys29-Pro528 is expressed with a 6His tag at the C-terminus.
Accession #	Q9NZU0
Host	Human Cells
Species	Human
Predicted Molecular Mass	57.6 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background



Alternative NamesLeucine-Rich Repeat Transmembrane Protein FLRT3; Fibronectin-Like Domain-
Containing Leucine-Rich Transmembrane Protein 3; FLRT3; KIAA1469BackgroundLeucine-Rich Repeat Transmembrane Protein FLRT3 (FLRT3) is a member of the
fibronectin leucine rich transmembrane protein (FLRT) family. Proteins in this family
play an role in cell adhesion and/or receptor signalling. FLRT3 is a single-pass type
I membrane protein and contains one fibronectin type-III domain, ten LRR
(leucine-rich) repeats, one LRRCT domain, and one LRRNT domain. FLRT3 may
have a function in cell adhesion and/or receptor signaling. FLRT3 may regulate
cellular adhesion between the epithelial apical ridge and the underlying
mesenchyme and in establishing the dorso-ventral position of the ridge.

Note

For Research Use Only, Not for Diagnostic Use.