


Product Name: Recombinant Human GITR (C-Fc)
Catalog #: PHH0731



Summary

Name	GITR/TNFRSF18/CD357/Tumor necrosis factor receptor superfamily member 18/Glucocorticoid-induced TNFR-related protein/AITR
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Glucocorticoid Induced TNF Receptor Family Related Gene is produced by our Mammalian expression system and the target gene encoding Gln26-Gln161 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q9Y5U5
Host	Human Cells
Species	Human
Predicted Molecular Mass	41.2 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human GITR (C-Fc)
Catalog #: PHH0731



Alternative Names

Tumor necrosis factor receptor superfamily member 18; TNFRSF18; Glucocorticoid-induced TNFR-related protein; CD357; TNFRSF18; AITR; GITR

Background

Tumor necrosis factor receptor superfamily member 18(Gitr) contains 3 TNFR-Cys repeats and it have four isforms.IsformA、 isformB and isformC is single-pass type I membrane protein and isformD is a secreted protein. The protein is the receptor for TNFSF18.It seems to be involved in interactions between activated T-lymphocytes and endothelial cells and in the regulation of T-cell receptor-mediated cell death. It mediated NF-kappa-B activation via the TRAF2/NIK pathway.It binds to TRAF1, TRAF2, and TRAF3, but not TRAF5 and TRAF6 and binds through its C-terminus to SIVA1/SIVA.It preferentially expressed in activated T lymphocytes and up-regulated in peripheral mononuclear cells after antigen stimulation/lymphocyte activation.

Note

For Research Use Only , Not for Diagnostic Use.