

Product Name: Recombinant Human GPD1 (C-6His)
Catalog #: PHH0755




Summary

Name	GPD1/Glycerol-3-phosphate dehydrogenase
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Glycerol-3-Phosphate Dehydrogenase [NAD(+)], Cytoplasmic is produced by our Mammalian expression system and the target gene encoding Met1-Met349 is expressed with a 6His tag at the C-terminus.
Accession #	P21695
Host	Human Cells
Species	Human
Predicted Molecular Mass	38.6 KDa
Formulation	Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 10% Glycerol, pH 8.0.
Shipping	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	

SDS-PAGE image

Product Name: Recombinant Human GPD1 (C-6His)
Catalog #: PHH0755



Alternative Names

Glycerol-3-Phosphate Dehydrogenase [NAD(+)] Cytoplasmic; GPD-C; GPDH-C; GPD1

Background

Glycerol-3-Phosphate Dehydrogenase [NAD(+), Cytoplasmic (GPDH-C)] belongs to the NAD-Dependent Glycerol-3-Phosphate Dehydrogenase family. GPDH-C plays a critical role in carbohydrate and lipid metabolism by catalyzing the reversible conversion of Dihydroxyacetone Phosphate (DHAP) and reducing Nicotine Adenine Dinucleotide (NADH) to Glycerol-3-Phosphate (G3P) and NAD⁺. GPDH-C is inhibited by zinc ions and sulfate. Mutations in this gene are a cause of transient infantile hypertriglyceridemia. GPDH-C is unlike Glyceraldehyde 3-Phosphate Dehydrogenase (GAPDH); they have different substrates.

Note

For Research Use Only , Not for Diagnostic Use.