

Product Name: Recombinant Human Haptoglobin (C-6His)
Catalog #: PHH0771




Summary

Name	Haptoglobin/HP/Zonulin
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Haptoglobin/Zonulin is produced by our Mammalian expression system and the target gene encoding Val19-Gln160(alpha chain)&Ile162-Asn406(beta chain) is expressed with a 6His tag at the C-terminus.
Accession #	P00738
Host	Human Cells
Species	Human
Predicted Molecular Mass	15.9&28.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human Haptoglobin (C-6His)
Catalog #: PHH0771



Alternative Names

Haptoglobin; Zonulin; HP

Background

Haptoglobin(HP) is a secreted protein which is a member of the peptidase S1 family of serine proteases. As a result of hemolysis, hemoglobin is found to accumulate in the kidney and is secreted in the urine. Haptoglobin captures, and combines with free plasma hemoglobin to allow hepatic recycling of heme iron and to prevent kidney damage. Haptoglobin also acts as an Antimicrobial; Antioxidant has antibacterial activity and plays a role in modulating many aspects of the acute phase response. Hemoglobin/haptoglobin complexes are rapidly cleared by the macrophage CD163 scavenger receptor expressed on the surface of liver Kupfer cells through an endocytic lysosomal degradation pathway. Uncleaved haptoglobin, also known as zonulin, plays a role in intestinal permeability, allowing intercellular tight junction disassembly, and controlling the equilibrium between tolerance and immunity to non-self antigens.

Note

For Research Use Only , Not for Diagnostic Use.