


Product Name: Recombinant Human HMGB2 (C-6His)
Catalog #: PHH0801



Summary

Name	HMGB2/High mobility group protein B2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human High Mobility Group Protein B2 is produced by our Mammalian expression system and the target gene encoding Gly2-Glu209 is expressed with a 6His tag at the C-terminus.
Accession #	P26583
Host	Human Cells
Species	Human
Predicted Molecular Mass	25.07 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 100mM NaCl, pH 8.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human HMGB2 (C-6His)
Catalog #: PHH0801



Alternative Names

High Mobility Group Protein B2; High Mobility Group Protein 2; HMG-2; HMGB2; HMG2

Background

High Mobility Group Protein B2 (HMGB2) belongs to the non-histone chromosomal high-mobility group protein family. Members of this family are chromatin-associated and widely spread in the nucleus of higher eukaryotic cells. HMGB2 contains 2 HMG box DNA-binding domains. It is associated with chromatin and has the ability to bend DNA, preferentially single-stranded DNA. It is shown that HMGB2 is able to efficiently bend DNA and form DNA circles. In addition, HMGB2 is involved in the final ligation step in DNA end-joining processes of DNA double-strand breaks repair and V(D)J recombination.

Note

For Research Use Only , Not for Diagnostic Use.