Product Name: Recombinant Human HMGB3 (C-6His)



Catalog #: PHH0802

Summary

Name HMGB3/High mobility group protein B3

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Human High Mobility Group Protein B3 is produced by our

Mammalian expression system and the target gene encoding Met1-Glu200 is

expressed with a 6His tag at the C-terminus.

Accession # 015347

Host **Human Cells**

Species Human

Predicted Molecular Mass 24 KDa

Lyophilized from a 0.2 µm filtered solution of 20mM PB, 8% Trehalose, 2% **Formulation**

Mannitol, 0.05% Tween80, 2mM EDTA, pH7.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Lyophilized protein should be stored at \leq -20°C, stable for one year after receipt. Stability&Storage


Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at \leq -20°C for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human HMGB3 (C-6His) Catalog #: PHH0802



Alternative Names

High Mobility Group Protein B3; High Mobility Group Protein 2a; HMG-2a; High

Mobility Group Protein 4; HMG-4; HMGB3; HMG2A; HMG4

Background

High Mobility Group Protein B3 (HMGB3) belongs to the HMGB family. Members of the HMG box subfamily are thought to be have an important role in DNA replication, nucleosome assembly and transcription. HMGB3 binds preferentiallly single-stranded DNA and unwinds double stranded DNA. HMGB3 consists of 200 amino acids and is localized to the cell nucleus. It contains two HMG box DNA-binding domain. HMGB3 binds preferentially single-stranded DNA and unwinds

double stranded DNA.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838