


Product Name: Recombinant Human HYOU1 (C-10His)
Catalog #: PHH0819



Summary

Name	Hypoxia up-regulated protein 1/HYOU1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Hypoxia Up-regulated Protein 1 is produced by our Mammalian expression system and the target gene encoding Met695-Leu999 is expressed with a 10His tag at the C-terminus.
Accession #	Q9Y4L1
Host	Human Cells
Species	Human
Predicted Molecular Mass	36 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human HYOU1 (C-10His)
Catalog #: PHH0819



Alternative Names

Hypoxia up-regulated protein 1; 150 kDa oxygen-regulated protein; ORP-150; 170 kDa glucose-regulated protein; GRP-170; HYOU1; ORP150

Background

Hypoxia up-regulated protein 1(HYOU1) is a member of the heat shock protein 70 family. Seven members from four different heat shock protein (HSP) families were identified including HYOU1, HSPC1(HSP86), HSPA5(Bip), HSPD1(HSP60), and several isoforms of the two testis-specific HSP70 chaperones HSPA2 and HSPA1L. HYOU1 is highly expressed in many tissues, such as liver, pancreas, macrophages within aortic atherosclerotic plaques, and in breast cancers. HYOU1 has a pivotal role in cytoprotective cellular mechanisms triggered by oxygen deprivation. It may play a role as a molecular chaperone and participate in protein folding. Suppression of HYOU1 is associated with accelerated apoptosis. It is suggested to have an important cytoprotective role in hypoxia-induced cellular perturbation. This protein has been shown to be up-regulated in tumors, especially in breast tumors, and thus it is associated with tumor invasiveness.

Note

For Research Use Only , Not for Diagnostic Use.