

Product Name: Recombinant Human IGFBP-5 (C-6His)
Catalog #: PHH0845




Summary

Name	IGFBP-5/Insulin-like Growth Factor-Binding Protein 5/IGFBP5
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Insulin-Like Growth Factor-Binding Protein 5 is produced by our Mammalian expression system and the target gene encoding Leu21-Glu272 is expressed with a 6His tag at the C-terminus.
Accession #	P24593
Host	Human Cells
Species	Human
Predicted Molecular Mass	29.61 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human IGFBP-5 (C-6His)
Catalog #: PHH0845



Alternative Names

Insulin-Like Growth Factor-Binding Protein 5; IBP-5; IGF-Binding Protein 5; IGFBP-5; IGFBP5; IBP5

Background

Insulin-Like Growth Factor-Binding Protein 5 (IGFBP-5) is a secreted protein that belongs to the insulin-like growth factor (IGF) binding proteins superfamily. Members of this family prolong the half-life of the IGFs. They have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors. IGFBP-5 contains one IGFBP N-terminal domain and one thyroglobulin type-1 domain. IGFBP-5 is expressed by fibroblasts, myoblasts and Osteosarcoma. It is also present at lower levels in liver, kidney, and brain.

Note

For Research Use Only , Not for Diagnostic Use.