## EnkiLife Product Name: Recombinant Human IL-18RAcP (C-6His) Catalog #: PHH0885

### **Summary**

Name IL-18RAcP/IL-18RAP/IL-18 Receptor Accessory Protein/IL-1R7

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/µg as determined by LAL test.

Construction Recombinant Human Interleukin-18 Receptor Accessory Protein is produced

by our Mammalian expression system and the target gene encoding Phe20-

Arg356 is expressed with a 6His tag at the C-terminus.

Accession # 095256

Host **Human Cells** 

**Species** Human

**Predicted Molecular Mass** 39.4 KDa

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. **Formulation** 

**Shipping** The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 Stability&Storage


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image



## **Background**

# Product Name: Recombinant Human IL-18RAcP (C-6His) Enkilife Catalog #: PHH0885

Alternative Names Interleukin-18 receptor accessory protein (IL18RAP); Accessory protein-like; CD218

antigen-like family member B; CDw218b; IL-1R accessory protein-like; Interleukin-1 receptor 7; Interleukin-18 receptor accessory protein-like; Interleukin-18 receptor

beta£"CD218b and IL1R7.

Background IL-18RAP is a single-pass type I membrane protein and contains two Iq-like C2-

type domains and one TIR domain, IL18RAP can be induced by IFN-alpha and IL12 in nature killer cells and T-cells. The coexpression of IL18R1 and IL18RAP is required for the activation of NF-kappa B and JNK in response to IL-18.Defects in

IL18RAP are associated with Coeliac disease.

#### Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838