


Product Name: Recombinant Human IL-1RAcP (Q9NPH3-2, C-6His)
Catalog #: PHH0892

Summary

Name	IL-1RAcP/IL1R3/IL-1 Receptor Accessory Protein (Q9NPH3-2,Ser21-Gln356)
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Interleukin-1 Receptor Accessory Protein is produced by our Mammalian expression system and the target gene encoding Ser21-Gln356 is expressed with a 6His tag at the C-terminus.
Accession #	Q9NPH3-2
Host	Human Cells
Species	Human
Predicted Molecular Mass	39.75 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human IL-1RAcP (Q9NPH3-2, C-6His)
Catalog #: PHH0892



Alternative Names

Interleukin-1 Receptor Accessory Protein; IL-1 Receptor Accessory Protein; IL-1RAcP; Interleukin-1 Receptor 3; IL-1R-3; IL-1R3; IL1RAP; C3orf13; IL1R3

Background

Interleukin-1 Receptor Accessory Protein (IL-1RAcP) is a member of the interleukin-1 receptor family. It contains three Ig-like C2-type domains in the extracellular region and a long cytoplasmic domain implicated in signal transduction. IL-1RAcP acts as a non-ligand binding accessory component of the receptors for IL1 α , IL1 β and IL33. IL-1RAcP mediates interleukin-1-dependent activation of NF-kappa-B. It is part of the membrane-bound form of the IL-1 receptor. IL-1 RAcP takes part in the Signaling ways by the formation of a ternary complex containing IL1R1, TOLLIP, MYD88, and IRAK1 or IRAK2. In addition, IL-1RAcP modulates the response to interleukins by associating with soluble IL1R1 and enhancing interleukin-binding to the decoy receptor.

Note

For Research Use Only , Not for Diagnostic Use.