Product Name: Recombinant Human IZUMO4 (C-Fc)

Catalog #: PHH1024



Summary

Name Izumo sperm-egg fusion protein 4/IZUMO4/C19orf36

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Izumo Sperm-egg Fusion Protein 4 is produced by our

Mammalian expression system and the target gene encoding His16-His214 is

expressed with a human IgG1 Fc tag at the C-terminus.

Accession # Q1ZYL8-2

Host Human Cells

Species Human

Predicted Molecular Mass 50.1 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human IZUMO4 (C-Fc)

Catalog #: PHH1024



Background

Alternative Names Sperm 22 kDa protein c113; IZUMO4; C19orf36; UNQ831/PRO1758

Background Izumo sperm-egg fusion protein 4 is a sperm membrane protein which plays a key

role in the fusion in the mouse. IZUMO4 has an N-terminal domain with significant homology to the N-terminal domain of Izumo.It belongs to the Izumo family . Izumo 4 is a soluble protein expressed in the testis and in other tissues. Izumo domain possesses the ability to form dimers, whereas the transmembrane domain or the cytoplasmic domain or both of Izumo 1 are required for the formation of

multimers of higher order.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838