## **Product Name: Recombinant Human SCF (C-6His)**

Catalog #: PHH1050



### **Summary**

Name SCF/Stem Cell Factor/c-kit Ligand/KITLG

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Human Stem Cell Factor is produced by our Mammalian

expression system and the target gene encoding Glu26-His214 is expressed

with a 6His tag at the C-terminus.

Accession # P21583

**Host** Human Cells

**Species** Human

Predicted Molecular Mass 22 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Lyophilized protein should be stored at  $\leq$  -20°C, stable for one year after receipt.


Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at  $\leq$  -20°C for 3 months.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

#### **SDS-PAGE** image



# **Product Name: Recombinant Human SCF (C-6His)**

Catalog #: PHH1050



### **Background**

Alternative Names Kit Ligand; Mast Cell Growth Factor; MGF; Stem Cell Factor; SCF; c-Kit ligand; KITLG;

MGF; SCF

Background Stem Cell Factor (SCF) is a hematopoietic growth factor that exerts its activity at

the early stages of hematopoiesis. SCF stimulates the proliferation of myeloid, erythroid, and lymphoid progenitors in bone marrow cultures and has been shown

to act synergistically with colony stimulating factors.

### Note

For Research Use Only , Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838