


Summary

| | |
|---------------------------------|--|
| Name | LAIR2/CD306 |
| Purity | Greater than 95% as determined by reducing SDS-PAGE |
| Endotoxin level | <1 EU/μg as determined by LAL test. |
| Construction | Recombinant Human Leukocyte-Associated Immunoglobulin-Like Receptor 2 is produced by our Mammalian expression system and the target gene encoding Gln22-Pro152 is expressed with a 6His tag at the C-terminus. |
| Accession # | Q6ISS4 |
| Host | Human Cells |
| Species | Human |
| Predicted Molecular Mass | 15.1 KDa |
| Formulation | Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| Stability&Storage | Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles. |
| Reconstitution | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

SDS-PAGE image

Product Name: Recombinant Human LAIR2 (C-6His)
Catalog #: PHH1062



Alternative Names

Leukocyte-Associated Immunoglobulin-Like Receptor 2; LAIR-2; CD306; LAIR2

Background

Leukocyte-Associated Immunoglobulin-Like Receptor 2 (LAIR2) is a secreted, 131 amino acid protein that contains one Ig-like C2 type domain, making it a member of the Ig superfamily. When compared to LAIR-1, its transmembrane counterpart, it shares 83% amino acid identity across the signal sequence and extracellular domains; although one is secreted and one is membrane-bound, the two LAIR proteins are thought to have arisen from a common gene ancestor and appear to share similar adhesion profiles. This suggests that LAIR-2 may compete with LAIR-1 for ligand binding. A 114 amino acid alternate splice form of LAIR-2 is truncated at the C terminus, but retains the entire Ig domain. The expression profile of these splice forms, and the presence of orthologs in other species, have not been reported.

Note

For Research Use Only , Not for Diagnostic Use.