

**Product Name: Recombinant Human LBP (C-6His)**  
**Catalog #: PHH1065**




---

## Summary

<b>Name</b>	LBP/Lipopolysaccharide-binding protein
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Lipopolysaccharide-Binding Protein is produced by our Mammalian expression system and the target gene encoding Ala26-Val481 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	P18428
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	51.95 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 500mM NaCl, 1mM EDTA, pH 8.0.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Human LBP (C-6His)**  
**Catalog #: PHH1065**



### **Alternative Names**

Lipopolysaccharide-Binding Protein; LBP

### **Background**

Lipopolysaccharide binding protein (LBP) is a plasma protein, belongs to a member of structurally and functionally related proteins which includes bactericidal permeability-increasing protein (BPI), plasma cholesteryl ester transfer protein (CETP) and phospholipid transfer protein (PLTP). It is involved in the acute-phase immunologic response to gram-negative bacterial infections. In cooperation with BPI, LBP binds LPS and interacts with the CD14 receptor, most likely playing a role in regulating LPS-dependent monocyte responses. Studies suggest that LBP is necessary for the rapid acute-phase response to LPS but not for the clearance of LPS from circulation. Finally, the LBP gene is found on chromosome 20, directly downstream of the BPI gene.

### **Note**

For Research Use Only , Not for Diagnostic Use.