


Product Name: Recombinant Human Lumican (C-6His)
Catalog #: PHH1108



Summary

Name	Lumican/LUM/LDC
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Lumican is produced by our Mammalian expression system and the target gene encoding Gln19-Asn338 is expressed with a 6His tag at the C-terminus.
Accession #	P51884
Host	Human Cells
Species	Human
Predicted Molecular Mass	37.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, 1mM EDTA, pH 7.2.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human Lumican (C-6His)
Catalog #: PHH1108



Alternative Names

Lumican; Keratan Sulfate Proteoglycan Lumican; KSPG Lumican; LUM; LDC; SLRR2D

Background

Lumican is a 40 kD secreted protein which belongs to the small leucine-rich repeat proteoglycans (SLRPs) and the class II subfamily. Human Lumican is synthesized as a 338 amino acid precursor then cut the 18 aa signal sequence. The mature Human Lumican contains 12 leucine-rich repeats (LRRs), 4 potential sites of N-linked glycosylation, and a C- terminal with two conserved cyst-eines. Lumican can be existed in extracellular matrix of human articular cartilage. Lumican participates in the maintenance of tissue homeostasis and regulates cellular functions in vivo, such as cell proliferation, adhesion, migration, and differentiation. The overexpression of lumican has been correlated to colorectal tumor, breast, neuroendocrine, and pancreatic cancers.

Note

For Research Use Only , Not for Diagnostic Use.