


**Product Name: Recombinant Human LYPD3 (C-6His)**  
**Catalog #: PHH1110**



## Summary

|                                 |  |
|---------------------------------|--|
| <b>Name</b>                     | Ly6/PLAUR domain-containing protein 3/C4.4A/LYPD3/MIG-C4   |
| <b>Purity</b>                   | Greater than 95% as determined by reducing SDS-PAGE  |
| <b>Endotoxin level</b>          | <1 EU/μg as determined by LAL test.  |
| <b>Construction</b>             | Recombinant Human Ly6/PLAUR Domain-Containing Protein 3 is produced by our Mammalian expression system and the target gene encoding Leu31-His286 is expressed with a 6His tag at the C-terminus.   |
| <b>Accession #</b>              | O95274   |
| <b>Host</b>                     | Human Cells  |
| <b>Species</b>                  | Human  |
| <b>Predicted Molecular Mass</b> | 27.89 KDa  |
| <b>Formulation</b>              | Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.  |
| <b>Shipping</b>                 | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.   |
| <b>Stability&amp;Storage</b>    | Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.   |
| <b>Reconstitution</b>           | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

## SDS-PAGE image



## Background

**Product Name: Recombinant Human LYPD3 (C-6His)**  
**Catalog #: PHH1110**

---



**Alternative Names**

Ly6/PLAUR Domain-Containing Protein 3; GPI-Anchored Metastasis-Associated Protein C4.4A Homolog; Matrigel-Induced Gene C4 Protein; MIG-C4; LYPD3; C4.4A

**Background**

Ly6/PLAUR domain containing3 (LYPD-3) is a GPI-linked protein. The structure of LYPD-3 is similar to the urokinasetype plasminogen activator receptor (uPAR). LYPD-3 is a 6 -100 kDa molecule with variable cell type-specific N-O-linked glycosylation, mature human LYPD-3 contains two uPAR/Ly6 domains and a Ser/Thr/Pro-rich (STP) region includes a protease sensitive site . The interaction of LYPD-3 with Laminin 1 and 5 on neighboring cells promotes the adhesion, spreading, and migration of tumor cells. LYPD-3 additionally interacts with Galectin-3 and the anterior gradient proteins AG-2 and AG-3. LYPD-3 overexpression in non-small cell lung cancer is predictive of increased mortality.

**Note**

For Research Use Only , Not for Diagnostic Use.