

Product Name: Recombinant Human MerTK (C-6His)
Catalog #: PHH1147




Summary

Name	MerTK/MERTK/MER/Receptor tyrosine kinase MerTK
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Tyrosine-protein Kinase Mer is produced by our Mammalian expression system and the target gene encoding Met1-Ala323 is expressed with a 6His tag at the C-terminus.
Accession #	Q1RMG3
Host	Human Cells
Species	Human
Predicted Molecular Mass	36 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human MerTK (C-6His)
Catalog #: PHH1147



Alternative Names

Tyrosine-protein kinase Mer; Proto-oncogene c-Mer; Receptor tyrosine kinase MerTK; MERTK; MER

Background

Tyrosine-protein kinase Mer (MERTK) is a single-pass type I membrane protein which belongs to the MER/AXL/TYRO3 receptor kinase family. MERTK include two fibronectin type-III domains, two Ig-like C2-type domains, and one tyrosine kinase domain. It can't be expressed in normal B- and T-lymphocytes, but it is usually expressed in numerous neoplastic B- and T-cell lines. MERTK could regulate many physiological processes, such as cell survival, migration, differentiation. It was demonstrated that the MERTK plays critical role in the engulfment and efficient clearance of apoptotic cells, platelet aggregation, and cytoskeleton reorganization. Not only these, it also plays an important role in inhibition of Toll-like receptors (TLRs)-mediated innate immune response by activating STAT1, which selectively induces production of suppressors of cytokine signaling SOCS1 and SOCS3. In addition, MERTK could regulate rod outer segments fragments phagocytosis in the retinal pigment epithelium (RPE), deficiency in MERTK are the cause of retinitis pigmentosa.

Note

For Research Use Only , Not for Diagnostic Use.