Product Name: Recombinant Human MMP-1 (C-6His) Catalog #: PHH1167

C EnkiLife

Summary

Name MMP-1/Interstitial collagenase

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Matrix Metalloproteinase-1 is produced by our

Mammalian expression system and the target gene encoding Phe20-Asn469 is expressed with a 6His tag at the C-terminus. The proenzyme needs to be

activated by APMA for an activated form.

Accession # P03956

Host Human Cells

Species Human

Predicted Molecular Mass 52.88 KDa

Formulation Supplied as a 0.2 µm filtered solution of 20mM MES, 150mM NaCl, 2mM CaCl2,

1mM DTT, 0.05%Brij35, 10% Glycerol, pH 5.0.

Shipping The product is shipped on dry ice/polar packs. Upon receipt, store it immediately

at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.


Reconstitution

SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human MMP-1 (C-6His) Catalog #: PHH1167

EnkiLife



Alternative Names

Interstitial Collagenase; Fibroblast Collagenase; Matrix Metalloproteinase-1; MMP-1; MMP1; CLG

Background

Matrix Metalloproteinase-1 (MMP-1) is expressed by fibroblasts, keratinocytes, endothelial cells, monocytes and macrophages. MMP1 contains several distinct domains: a prodomain that is cleaved upon activation, a catalytic domain containing the zinc binding site, a short hinge region, and a carboxyl terminal (hemopexin like) domain. MMP-1 can degrade a broad range of substrates including types I, II, III, VII, VIII, and X collagens as well as casein, gelatin, α1 antitrypsin, myelin basic protein, L-Selectin, pro-TNF, IL1, IGFBP3, IGFBP5, pro-MMP2, and pro-MMP9. A significant role of MMP1 is the degradation of fibrillar collagens in extracellular matrix remodeling, characterized by the cleavage of the interstitial collagen triple helix into 3/4, 1/4 fragments. MMP1 may also be involved in enzyme cascades, cytokine regulation and cell surface molecule modulation.

Note

For Research Use Only, Not for Diagnostic Use.