


Summary

Name	NKp30/CD337/NCR3
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Natural Cytotoxicity Triggering Receptor 3 is produced by our Mammalian expression system and the target gene encoding Leu19-Thr138 is expressed with a 6His tag at the C-terminus.
Accession #	O14931
Host	Human Cells
Species	Human
Predicted Molecular Mass	12.84 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human NCR3 (C-6His)
Catalog #: PHH1199



Alternative Names

NCR3 Protein; Natural Cytotoxicity Triggering Receptor 3; Natural Cytotoxicity Triggering Receptor 3 Isoform CRA_c; NCR3

Background

Natural Cytotoxicity Triggering Receptor 3 (NCR3) along with Nkp44 and Nkp46 constitute a group of receptors termed "Natural Cytotoxicity Receptors". They play a major role in triggering NK-mediated killing of most tumor cells lines. Nkp30 is a type I transmembrane protein having a single extracellular V-like immunoglobulin domain. Nkp30 is selectively expressed both in resting and activated human NK cells. In addition, Nkp30 is also involved in NK-mediated induction of dendritic cell (DC) maturation. It has been demonstrated that NK cell activation signaling specifically induces lytic activity against several tumor cell types and synthesis of new NF- κ B dependent proteins during the initiation of cytotoxicity.

Note

For Research Use Only , Not for Diagnostic Use.