## **Summary**

Name NKG2D Ligand 2/NKG2DL2/N2DL2/ULBP-2

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Human UL16 Binding Protein-2/NKG2D ligand 2 is produced by

our Mammalian expression system and the target gene encoding Gly26-

Ser217 is expressed with a 6His tag at the C-terminus.

Accession # Q9BZM5

**Host** Human Cells

**Species** Human

Predicted Molecular Mass 22.78 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image



# **Background**

# Product Name: Recombinant Human NKG2DL2 (C-6His) Enkilife Catalog #: PHH1233

Alternative Names NKG2D Ligand 2; N2DL-2; NKG2DL2; ALCAN-Alpha; Retinoic Acid Early Transcript

1H; UL16-Binding Protein 2; ULBP2; N2DL2; RAET1H

**Background** NKG2D Ligand 2 (N2DL2) is a member of a family of cell-surface proteins. N2DL2

function as ligands for human cytomegalovirus glycoprotein UL16. N2DL2 is anchored to the membrane via a GPI-linkage. N2DL2 is bind to human NKG2D, an activating receptor expressed on NK cells, NKT cells, T cells. Engagement of NKG2D results in the activation of cytolytic activity and cytokine production by these effects cells. The ULBPs are expressed on some tumor cells and have been

implicated in tumor surveillance.

#### Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838