Product Name: Recombinant Human NPTX1 (C-6His)

Catalog #: PHH1240



Summary

Name NPTX1

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Neuronal Pentraxin-1 is produced by our Mammalian

expression system and the target gene encoding Gln23-Asn432 is expressed

with a 6His tag at the C-terminus.

Accession # Q15818

Host Human Cells

Species Human

Predicted Molecular Mass 45.9 KDa

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, 1mM EDTA, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is


not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human NPTX1 (C-6His)

Catalog #: PHH1240





Alternative Names

Neuronal pentraxin-1;NPTX1;NP1

Background

Neuronal Pentraxin (NPTX1, NP1) is a secreted glycoprotein within the Pentraxin family. NPTX1 is co-expressed and forms heteromultimers with the related secreted protein, NPTX2/NARP, NPTXR (Neuronal Pentraxin Receptor) at excitatory synapses. Mature human NPTX1 shares 97% aa sequence identity with mouse, and rat NPTX1. It is produced by hippocampal, cerebral and cerebellar neurons, retinal ganglia and the inner nuclear layer of the retina. It is enriched on presynaptic axonal membranes where it forms complexes with NPTXR. It may be involved in mediating uptake of synaptic material during synapse remodeling or in mediating the synaptic clustering of AMPA glutamate receptors at a subset of excitatory synapses.

Note

For Research Use Only , Not for Diagnostic Use.